

**Application
for
United States Letters Patent**

09975607-10101
TOTOT 20952660

To all whom it may concern:

Be it known that we

*Kathryn S.E. Cheah and Kenneth K.M. Cheung
have invented certain new and useful improvements in*

USES OF TRANSGENIC ANIMALS CONTAINING A TYPE X COLLAGEN MUTANT

of which the following is a full, clear and exact description.

**USES OF TRANSGENIC ANIMALS CONTAINING
A TYPE X COLLAGEN MUTANT**

This application claims the benefit of U.S. Provisional Application No. 60/103,550, filed October 8, 1998, the contents of which are hereby incorporated by reference.

5 Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

10

Background of the Invention

Type X Collagen and its function

15 Type X collagen is a homotrimer of three $\alpha 1(X)$ chains, with a short (38aa) non-helical amino terminus (NC2), a triple helix of 463aa and a C-terminal highly conserved noncollagenous domain (NC1) of 161aa. This collagen is the major extracellular component synthesized by hypertrophic chondrocytes in growth cartilage destined to be calcified and in zones of secondary ossification
20 (1,2). Expression of the $\alpha 1(X)$ collagen gene is specifically associated with hypertrophic chondrocytes and precedes the onset of endochondral ossification (3). Although this collagen does not form fibrils, it has been found as fine pericellular filaments in association with
25 cartilage collagen fibrils (4). Type X collagen molecules may also form other supramolecular structures in the matrix, since they have been shown to assemble into a hexagonal lattice in vitro (5).

30 Apart from association with collagen fibrils, type X collagen interacts with other matrix components, such as annexin V, chondrocalcin (6) and proteoglycans (4). Type X collagen has also been shown to be intimately associated with the calcification process by binding to
35 Ca^{++} and matrix vesicles which are cell-derived

09975607 101104

microstructures found in the matrix of calcifying cartilage and bone and thought to be important in the initiation of mineral deposition (7). In addition expression of type X collagen precedes mineral deposition by cultured chondrocytes (8).

Despite the wealth of information about type X collagen, the precise function of this protein and its role in the pathogenesis of chondrodysplasia, has remained the subject of controversy. Because of its specific association with hypertrophic chondrocytes in the calcifying zone of growth plate cartilage, type X collagen has been proposed to be important for endochondral bone formation (2). Proposed functions include, providing an easily resorbed fabric for the deposition of bone matrix during endochondral growth of long bones; providing support as the cartilage matrix is degraded during endochondral ossification (9,10); or regulating the calcification process during endochondral ossification (11-14). Reconciling these opposing views has also been difficult because the consequences of gene mutations which result in type X collagen deficiency in human and mouse differ.

Mutations in the NC1 encoding domain of the human $\alpha 1(X)$ collagen gene (COL10A1) have been found to be associated with the autosomal dominant inherited skeletal disorder, Schmid metaphyseal chondrodysplasia (SMCD) (15-18). SMCD is a relatively mild form of metaphyseal chondrodysplasia, resulting from growth plate abnormalities. The SMCD phenotype is variable in severity and characterized by short to normal stature, with genu varum (bow legs), coxa vara (a reduced angle between the femoral neck and shaft) and flaring of the metaphyses of long bone (19,20). Transgenic mice expressing truncated chicken type X collagen, display much more severe skeletal abnormalities, similar to human

09075607-101101

spondylometaphyseal dysplasia(SMD) (21) in which there is compression of the hypertrophic zone of the growth plate and a decrease in newly formed bony trabeculae.

5 The phenotypes of SMCD patients and the SMD-like transgenic mice favour a supportive role for type X collagen (15,16,22). Therefore it was surprising to find that mice carrying a null mutation in the $\alpha 1(X)$ collagen gene (Coll10a1) have been reported to show no abnormality
10 and no signs of SMCD (23). K. Cheah in collaboration with others had also created a null mutation in mouse Coll10a1 by homologous recombination in ES cells, to gain insight into the function of type X collagen (24). To resolve the apparently contradictory consequences of
15 mutations in the gene in human and mouse and gain better insight into the pathogenesis of SMCD, we focused our study on the consequences of type X collagen deficiency on the structure of the growth plate and trabecular bone, and on the organization of matrix components within
20 cartilage.

This disclosure showed that type X collagen deficiency in mice does have phenotypic consequences which partly resemble SMCD, reducing the apparent discrepancy in
25 phenotype between human and mouse (24). Intriguingly, the major impact of type X collagen deficiency does not lie in its site of expression, the hypertrophic zone, but rather affects other zones of the growth plate and in bone. It is also found that the consequence of loss of
30 type X collagen in mutant mice is a major change in the distribution of matrix materials such as proteoglycans and matrix vesicles, within the epiphyseal cartilage. Other features of type X collagen deficiency are a significant compression of the resting zone and articular
35 cartilage.

Phenotypic features which partly resemble SMCD were

00975607-101101
TOTOT 20952660

found, such as persistence of cartilage in trabecular bone, alterations in bone mineralization and trabecular structure. In particular, type X collagen deficient mice develop coxa vara, one of phenotypic changes common in human SMCD. These findings have led us to propose a function for type X collagen. Based on these findings we propose that type X collagen plays a role in the normal compartmentalization of the cartilage matrix. Type X collagen deficiency alters the distribution of cartilage matrix components thereby impacting on the supporting properties of the growth plate and the mineralization process, resulting in abnormal trabecular bone. This hypothesis would accommodate the previously conflicting views of the function of type X collagen and of the molecular pathogenesis of SMCD.

0995607 10101
TOTTOT 20952660

Summary of the Invention

This invention provides an isolated DNA comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth. In an embodiment, the DNA comprises the sequence of Coll10-13del as set forth in Figure 2. This invention also provides a vector which comprises the above-described DNA.

10 This invention also provides a method for production of the a polypeptide which regulates bone growth comprising a host-vector system which comprise the above vector and an appropriate host.

15 This invention provides a polypeptide encoded by the isolated DNA comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth. This invention also provides a polypeptide which comprise the portion of the mutated collagen X capable of regulating bone growth.

20 This invention provides a composition comprising the above polypeptide and a suitable carrier. This invention provides a pharmaceutical composition for increasing bone growth comprising the above polypeptide and a pharmaceutically acceptable carrier.

25 This invention also provides a method of treating a subject afflicted with dwarfism comprising administering to the subject an amount of the polypeptide comprising a portion of the mutated collagen X capable of regulating bone growth or the isolated DNA comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth effective to reverse the dwarfism.

09975607-101101

5 This invention provides a method of treating a subject afflicted with low bone mass comprising administering to the subject an amount of the polypeptide comprising a portion of the mutated collagen X capable of regulating bone growth or the isolated DNA comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth effective to treat low bone mass in the subject.

10 This invention provides a method of improving the quality and speed of bone union after fracture in a subject comprising administering to the subject an amount of the polypeptide comprising a portion of the mutated collagen X capable of regulating bone growth or the isolated DNA
15 comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth effective to improve the quality and speed of bone union.

20 This invention further provides a transgenic animal comprising an isolated DNA comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth.

09975607-101101
TOTOT 20952660

Brief Description of the Figures

Fig. 1. Diagram showing the structure of the Coll10-13del transgene. Collagen X gene and protein structure.

5

Fig. 2. Comparison of the mouse and human nonsense sequence generated from an equivalent 13 base pairs deletion within the NC1 domain of collagen X. Amino acids are numbered from the start of translation (3). The underlined bases in the wt sequence (SEQ ID NO: 1) represent the 13 bp deleted in the mutant sequence (SEQ ID NO: 2). Variations between the mouse and human nonsense sequences are highlighted in bold. The boxed amino acids represent a mutant specific polypeptide synthesized for polyclonal antibody production in rabbit. The nonsense amino acid sequences resulting from the deletion in the mouse and human are aligned for comparison. Amino acids are numbered from the start of translation(3). The underlined bases in the wt sequence represent the 13 bp deleted in the mutant sequence. Variations between the mouse and human nonsense sequences are highlighted in yellow. The boxed amino acids represent a mutant specific polypeptide synthesized for polyclonal antibody production in rabbit. Wild type amino acid sequence (SEQ ID NO: 3). Mutant mouse amino acid sequence (SEQ ID NO: 4). Mutant human amino acid sequence (SEQ ID NO: 5).

10

15

20

25

30

Fig. 3. Pictures of 6 month old Coll10-13del transgenic (Tg) and non-transgenic (NT) mice and X-rays. X-rays shown here and in subsequent figures are taken at the same exposures using a mammogram machine. A and B the scale bar represents 1cm;

35

00075607-101101

C-E scale bar represents 0.5cm. A. Left 34.10.57(NT), right 34.10.54 (Tg) at 6 months of age showing differences in overall body length and shape, in particular differences in the length of the limbs. B. Left 34.10.57(NT), right 34.10.224(Tg) at 6 months, showing differences in the shape of the head. C-E. 34.10.372 (NT), 34.10.373 (Tg) and 34.10.224 (Tg) respectively. Demonstrates the variability in severity of the malformation of the feet and digits. E also has an extra digit. F. X-ray of 34.10.224, showing the presence of an extra digit.

Fig. 4. X-ray images of 5 week old Col10-13del mice compared to non-transgenic mice. Note marked hyperostosis of bones as indicated by the increased opacity (whiteness) of image. A-B. 34.10.372(NT) and 34.10.373(Tg) top views. C-D 34.10.372 and 34.10.373 lateral view.

Fig. 5. X-rays of 10 week old Col10-13del mice compared to non-transgenic mice. A-B. 34.10.57(NT) and 34.10.54(Tg) top views. C-D 34.10.57 and 34.10.54 lateral view.

Fig. 6. X-rays of 15 week old Col10-13del mice compared to non-transgenic mice. A-B. 34.10.57(NT) and 34.10.54(Tg) top views. C-D 34.10.57 and 34.10.54 lateral view.

Fig. 7. Histology of growth plates of Col10-31del and wild-type littermate. A, C, E are wild-type non-transgenic (NT); B, D, F are Col10-13del transgenic (T). Note the greatly increased heights of the proliferating (p) and hypertrophic (h) zones in the Col10-13del mice

09975607-101101

as compared to wild-type littermate.

Fig. 8. Expression of p57kip2, PCNA and Hsp47 in 2 days post partum distal femur growth plates of Coll10-13del transgenic and non-transgenic mice. Expression shown are: a,b) cyclin-dependent kinase inhibitor p57kip2; c,d) proliferating cell nuclear antigen (PCNA, antibody from Santa Cruz Biotechnology, Santa Cruz, California); e,f) heat shock protein 47 (Hsp47). Positive signal in cells is shown by the brown immunoperoxidase staining using Vectastain ABC kit (Vector Laboratories, Burlingame, California). Staining of trabecular bony matrix is non-specific. All images are at 100X magnification except boxes are regions magnified 200 times. T: trabecular bone, H: hypertrophic zone. P: proliferative zone.

Fig. 9. Expression of wild type and mutant collagen X. Immunofluorescence (FITC staining) on cryosections from new born distal femur growth plates of Coll10-13del transgenic (tg, a,c,d) and non-transgenic mice (nt,b). a,b) ColX antibody staining; c) 13-del mutant protein antibody staining. Note punctate staining in the 13 del antibody stained section from transgenic growth plat, indicating mainly intracellular staining of hypertrophic chondrocytes (c) compared to extracellular staining with collagen X antibody staining of non-transgenic sections (b). In non-transgenic sections, only very weak background was seen (not shown). d) Immunostaining showing intracellular localisation of the Coll10-13del protein in osteoblast-like cells found in the trabecular bone in (c). T: trabecular bone H:

hypertrophic zone.

09975607.101101

This invention provides an isolated DNA comprising the sequence which codes for a mutated collagen X or a portion thereof wherein the expression of said DNA regulates bone growth. It is the intention of this invention to cover all mutant collagen X which is capable to regulate bone growth. This invention also covers portion of mutated collagen X which has the biological activities of regulating bone growth and polypeptides which bears this domain (portion).

This invention also provides a vector which comprises the above-described DNA.

Vectors which comprise the isolated nucleic acid molecule described hereinabove also are provided. Suitable vectors comprise, but are not limited to, a plasmid or a virus. These vectors may be transformed into a suitable host cell to form a host cell vector system for the production of a polypeptide capable of regulating bone growth.

This invention further provides an isolated DNA or cDNA molecule described hereinabove wherein the host cell is prokaryote or eukaryote. For example, the cells may be bacterial cells (such as E.coli), yeast cells, fungal cells, insect cells or animal cells. Suitable animal cells include, but are not limited to Vero cells, HeLa cells, Cos cells, CV1 cells and various primary mammalian cells.

This invention also provides a polypeptide encoded by

above-described DNA. This invention also provides polypeptide which bears the domain(s) of the mutant collagen X which regulates bone growth.

5 This invention also provide a composition comprising the above-described polypeptide and a suitable carrier. A suitable carrier are carrier which is capable of dissolving the polypeptide and yet not affect the biological activity of said polypeptide. For example,
10 the carrier may be a physiologically saline.

This invention also provides a pharmaceutical composition for increasing bone growth comprising the above-described polypeptide and a pharmaceutically acceptable carrier.

15 For the purposes of this invention "pharmaceutically acceptable carriers" means any of the standard pharmaceutical carriers. Examples of suitable carriers are well known in the art and may include, but not
20 limited to, any of the standard pharmaceutical carriers such as physiologic saline solution, phosphate buffered saline solutions, phosphate buffered saline containing Polysorb 80, water, emulsions such as oil/water emulsion, and various type of wetting agents. Other carriers may
25 also include sterile solutions, tablets, coated tablets, and capsules.

Typically such carriers contain excipients such as starch, milk, sugar, certain types of clay, gelatin,
30 stearic acid or salts thereof, magnesium or calcium stearate, talc, vegetable fats or oils, gums, glycols, or other known excipients. Such carriers may also include flavor and color additives or other ingredients. Compositions comprising such carriers are formulated by
35 well known conventional methods.

This invention also provides a method of treating a

009560 10101

subject afflicted with dwarfism comprising administering to the subject an amount of the above polypeptide or the above-described DNA of effective to reverse the dwarfism.

5 Methods of determining an "effective amount" are well known to those skilled in the art and depend upon factors including, but not limited to: the size of the patient and the carrier used.

10 This invention provides a method of treating a subject afflicted with low bone mass, including but not limited to osteoporosis, comprising administering to the subject an amount of the above polypeptide or DNA via for example the administration of bone marrow/stromal cells carrying
15 and expressing the Col10-13del transgene.

20 This invention provides a method of enhancing the quality of fracture healing and of the healed bone of a subject comprising administering to or implanting into or around the bone of the subject an amount of the above polypeptide or DNA comprising the above-described DNA effective to enhance the quality of fracture healing and of the healed bone.

25 As used herein, enhancing the quality of fracture healing means an increase in the rate of callus and bone formation, a reduction in the incidence of failure to heal (non-union), an increase in the quantity of bone formed, or an improvement in the mechanical properties of
30 the callus or of the bone formed. Enhancement of the healed bone means an improvement in the mechanical properties of the bone or and increase in the quantity of bone formed.

35 This invention also provides to a patient with low bone mass to improve the quality of the healed bone. Since type X collagen protein is an extracellular matrix

09975607-101101

component, it and its mutant forms may be administered extracellularly via local therapy which involves biodegradable vehicles. Methods of administering the polypeptide includes implanting coated or impregnated forms of solid support material e.g. ceramic powders (Gordon, E., Lasserre, A, Stull, P., Bajpai, P.K., England, B. 1997 Biomed Sci Instrum 33: 131-136); polymethymethacrylate cement; biodegradable polymers (such as poly lactides-co-glycolides Ramchandani M. Robinson, D. 1998 J. Controlled Release 54: 167-175), collagen vehicles such as resorbable collagen membranes (King, G.N., King, N., Hughes, F.J. 1998. J. Periodontal Res 33: 226-236) or gels impregnated with extracellular matrix proteins and the polypeptide and other similar approaches.

Methods of allowing site-specific gene delivery of the DNA may include implanting osteoblasts which have been transfected with the Col10-13del transgene and which express the transgene. We have previously identified a subset of the regulatory sequences in the Col10a1 gene (within the whole col10-13del construct) which can direct expression specifically to endosteal cells and osteoblasts (Cheah and Poon, unpublished results). Alternatively, implants containing chondrocytes transfected with the Col10-13del construct and expressing the transgene may be transferred to bone defects. Liposome based vehicles may also be used as a means to deliver the Col10-13del transgene to cells at the fracture site.

In an embodiment, the above-described DNA is operatively linked to inducible regulatory element.

This invention further provides a transgenic animal comprising an the isolated DNA comprising the sequence Col10-13del as set forth in Figure 2 or a portion thereof

00075607 101101

of which regulates bone growth when expressed.

In an embodiment, the transgenic animal of claim is a mouse.

5

This invention also provides a transgenic animal comprising an DNA designated Col10-13del as set forth in Figure 2.

10

This invention provides a method for identifying whether an agent which stimulates bone growth comprising steps of:

15

a) administering the agent to the above transgenic animal; and b) examining the transgenic animal after the administration of the agent to determine whether bone growth has been stimulated.

20

This invention also provides a method for assessing the effect of surgery on increasing bone length comprising steps of: a) performing surgery on the above transgenic animal; and b) assessing the increase in bone length of the animal to determine the effect of surgery on said animal.

25

In an embodiment, the assessment is performed by direct measurement of the increase in bone length and the animal is sacrificed at different time for the accurate measurement.

30

The invention also provides a method for assessing the effect of bone growth stimulating agents on fracture repair comprising steps of: a) creating a fracture repair in the above-described transgenic animal; b) administering the agent to the transgenic animal; and b) examining the transgenic animal after the administration of the agent to determine whether bone growth has been stimulated.

35

09975607-101101

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

09975607 101101
TOTTOT 20952660

Experimental Details

Molecular mechanisms underlying SMCD type X collagen mutations in humans

5 An important issue raised by our findings, is the
relative mildness of phenotype in the null mutants
compared with the SMD-like transgenic mice and human SMCD
(reviewed in 25). Since collagen α chains associate via
the NC1 domain, it has been proposed that in human SMCD,
10 the NC1 mutations result in failure of trimer assembly
(15,16) and the phenotypic changes are caused because
type X collagen is depleted in the matrix. This proposal
is supported by recent reports that collagen α 1(X) chains
carrying SMCD mutations are unable to form trimers in in
15 vitro assembly and cell transfection experiments (26,
27). In the SMD transgenic mice, the chondrodysplasia is
postulated to be caused by a depletion of type X collagen
as a result of instability/degradation of mutant
chicken-mouse hybrid protein (21) or because the chicken
20 α chains interfere with the normal assembly of mouse type
X collagen. The ability of α 1(X) chains with internal
deletions within the helix to assemble with normal chains
as heterotrimers and be secreted, support the latter
explanation (27). It is notable that the changes in
25 trabecular structure observed in SMD-like transgenic mice
(21) were similar but more severe than in the type X
collagen null mutants. Our results suggested that
different phenotypes of the null mutants and the SMD-like
transgenic mice probably reflect differing severity of
30 outcome of mutations resulting in loss versus gain of
function.

The reason for the late onset of coxa vara and the
relatively milder phenotypic changes in type X collagen
35 mutant mice, compared with human SMCD, is still not fully
understood. The late onset of the defect in the mutant
mice compared with the SMCD patients may be related to

00075607 101101

the differences in loading of the growth plate between man, a biped and mouse, a quadriped. However, the type X collagen content in SMCD growth plates has not been determined. In addition it is not known if synthesizing abnormal $\alpha 1(X)$ collagen chains which cannot assemble, has additional effects on hypertrophic chondrocytes, affecting their growth. It would therefore be important to compare the phenotypic consequences of the null with a SMCD mutation in mice.

Experimental Results & Discussion

Expressing type X collagen chains containing a SMCD mutation in transgenic mice

We have identified genomic sequences in the 5' flanking region, first and second introns of the mouse Col10a1 gene which were sufficient to drive tissue-specific expression of a FLAG- epitope-marked form of collagen X in hypertrophic chondrocytes in transgenic mice. There was also some expression in some chondrocytes in the lower proliferative zone. Expression was also seen in a few cells in the bone marrow compartment which could be osteoblasts. We used these sequences to initiate experiments designed to address these questions and have started to study the phenotypic consequence of expressing collagen $\alpha 1(X)$ chains carrying a SMCD mutation in hypertrophic chondrocytes of mice heterozygous and homozygous null for type X collagen.

Sited-directed mutagenesis (15) is used to introduce the mutation into a genomic clone containing the whole mouse collagen X gene (Col10a1) and the resulting clone is designated Col10-13del. The mutation was a 13-bp deletion within the conserved NC1 domain (Fig. 1). This mutation causes a shift in the amino acid reading frame at position 619 resulting in the production of a nonsense sequence (52 amino acids) from residue 620 with a

00975607-101101

prediction premature termination at 671, 9 amino acids shorter than the wild type (Fig. 2). This mutant construct, Coll10-13del was introduced into the germ-line of mice wild-type or heterozygous for Coll10a1, by transgenesis. Three transgenic mouse lines carrying this transgene on a wild-type Coll10a1 genetic background were found to have abnormal skeletal phenotype which, although variable in severity, was very similar in these lines. All three of these lines expressed the transgene as assessed by RT-PCR on RNAs from cartilage of the mice. Surprisingly the external gross phenotype of these transgenic mice was not similar to human SMCD but rather resembled other human skeletal disorders as follows:

Gross phenotype: The phenotype observed in the Coll10-13del mice appeared to be a rhizomelic skeletal dysplasia with analogy to human cranio-tubular remodelling disorders. The onset of observable differences in phenotype between transgenic and wild-type mice occurs 4-6 days post-partum. The transgenic mice are smaller than their littermates. The observed phenotype at this stage involves the distal digits. The digits are shorter and fatter than their wild type counterparts and appear to be hypermobile (Fig. 3). There is variation in degree in which the digits are short, bent or curved outwards, but this is a common feature of all the transgenic mice. Fore- and hind-limbs are affected, often the latter more severely. As the mice develop, a rounding of the fronto-nasal area becomes apparent. X-ray analysis of the mice from 5 weeks (5,10,15 wks) onwards has revealed a progressive hyperostosis (hyper =3D increased, ost =3D bone, osis =3D condition i.e. a condition of increased bone), which affects skull, ribs, spine and long bones (Figs. 4-6). As with the distal digits the long bones (humeri and femurs), ribs and pelvic elements are also shorter and broader than those observed in the wild type mice. There

09975607 101101

is also a strange lucency and expansion of the medullary cavity distal to the mid-point of the femurs. By 10-15 weeks the mice show varying degrees of waddling gait with splayed feet, with those most severely affected showing a "dragging posture" of their rear end probably because the hind legs are not normally flexed.

Histopathology: Comparison of the histology of sections from the growth plates of 10day and 4 week non-transgenic and Coll10-13del mice show marked differences (Fig 7). The proliferative zone of transgenic mice is enlarged compared to non-transgenic littermates. The most dramatic changes are seen in the hypertrophic zone which is considerably increased in height in Coll10-13del mice, with hypertrophic chondrocytes appearing disorganized and reduced in size. In addition, the trabeculae at the chondro-osseous region in mutants appeared increased in size (Fig7).

Ultrastructure: ultrastructural changes in the growth plates of the Coll10-13del mice were analyzed. Initial analyses of 10day neonates show that the matrix organization in the transgenic mice within the proliferating zone is such that a distinct pericellular matrix compartment is missing, whereas in the non-transgenic it is present. In the hypertrophic zone this compartment is present in both groups. The density of organelles within the cell cytoplasm (rough endoplasmatic reticulum, Golgi, mitochondria, etc.) within the cells of transgenic mice is in the proliferating zone significantly reduced, whereas in the non-transgenic proliferating cells it is of a normal density. In the hypertrophic zones this value is much more difficult to assess; however, in the transgenic mice we identified an increased density of extended and filled endoplasmatic reticulum membrane cisterns, which are not present in non-transgenic mice. The significance of

09975607-101101

these findings will need further analyses. However it is possible that it may be that the ultrastructural changes are the result of two effects: in the proliferating zone, which lacks type X collagen, ectopic expression of the coll10-13del transgene causes interference/alteration in matrix assembly and affects chondrocyte proliferation. In the hypertrophic zone where the normal type X collagen is also present, mutant-wild-type heterotrimers may be forming which are difficult to secrete or are targeted for degradation, causing distension of the ER.

Molecular bases of the growth plate and bone abnormalities-

- 15 The occurrence of hyperostosis in these mice is an interesting phenomenon, especially since type X collagen is not normally synthesized in bone. There is however some evidence that residual collagen X is present in the cartilaginous remnants within bone trabeculae (24).
- 20 There are several possible underlying causes, which singly or in combination, may underlie the abnormal bone phenotype in these mice:
- a. In both the human and mouse collagen X, the mutation results in a similar nonsense amino acid sequence terminating at the same residue. However, 15 of the 52 nonsense amino acids are different between the mouse and human sequence (Fig. 2) . This represents 29% variation and it is possible that the altered peptide sequence may contribute towards the phenotypic changes for example behaving as a signalling molecule/growth factor.
 - b. The phenotypic changes could be the result of ectopic expression of the transgene in the lower proliferative zone and/or osteoblasts
 - c. Synthesis of abnormal collagen X molecules which form heterotrimers with wild type chains
 - d. The phenotype is caused by the increased degradation

of mutant molecules by the hypertrophic chondrocytes.

e. Since collagen X has been shown to have a strong affinity for proteoglycans, the bone overgrowth could be caused by an imbalance in the matrix components, especially proteoglycans. This alteration in proteoglycan deposition or biosynthesis could contribute to the bone overgrowth since these molecules have been shown to be important for bone growth and remodelling.

f. The consequence of expressing abnormal collagen X could be to disrupt its interaction intracellularly with chaperones and extracellularly with other matrix molecules, proteinases and/or receptors (e.g integrins). These altered interactions are likely to alter intracellular signal transduction and hence chondrocyte proliferation and/or differentiation. The impact of such alterations are likely to be mostly at the chondro-osseous junction.

Further experimentation:

A. Basic Studies

Studying the molecular basis of the abnormalities in these mice will help us to understand the relationship between genotype and phenotype and will contribute to a further definition of the role of type X collagen in the formation and structural integrity of growth cartilage. This insight will be extremely valuable for an understanding of the pathology of diseases which result in the degeneration of cartilage, such as osteoarthritis. The progressive increase in bone density in these mice also imply deregulation of bone growth and remodelling. It is not yet clear if the phenotypic abnormalities are a primary or secondary effect of expression of the Col10-13del transgene. However this abnormality raises many questions about the regulatory pathways for bone growth. Therefore these mice will be used to:

00975607 10101

- a) identify key molecules and pathways important for the regulation of bone growth, with implications for human bone disorders.
- b) study differentiation, proliferation, apoptosis and cell-cycle regulation in the growth plate.
- c) study the role of the marrow microenvironment and the chondro-osseous junction on hematopoiesis, vascularization of bone, bone formation and remodelling.

The following studies will be performed:

- i) Phenotype and molecular characterization

Cartilage, the growth plate and membranous bone of mutants will be extensively analyzed. The spatial and temporal expression pattern of the transgene during development and postnatal growth these mice will be studied further by insitu hybridization and immunohistochemistry. In situ hybridization analyses will be used to determine if the transgene is expressed at the appropriate time in development as the wild type gene. Specific oligonucleotide or riboprobes will be generated spanning the deletion site. Conditions of hybridization will be optimized to give specific hybridization only to the transgene mRNA. In addition since our previous studies with the wild type gene marked by a FLAG epitope had shown expression in the hypertrophic chondrocytes as well as some proliferating chondrocytes. In situ hybridization using wild type Col10a1 probe will show the overall distribution of both mutant and normal transcripts and assess if the Col10-13del transgene shows similar ectopic expression. Since the 13bp deletion creates a new peptide sequence in the NC1 domain, antibodies can be raised against the altered peptide and used for immunohistochemistry. This will enable us to determine the localization of the mutant collagen X protein. Attention will also be taken to determine if mutant protein is present in the bone in

the cartilaginous remnants within boney trabeculae.

Levels of expression of the mutant transcript relative to wild-type will be determined using RNase protection assays using a riboprobe spanning the mutation, enabling both type of transcripts to be detected in one assay.

Proper chondrocyte differentiation will be assessed using appropriate molecular markers (30, 31), including type IIA procollagen (marker for pre-chondrocytes); long-form Col9a1, Col11a2, aggrecan (expressed by proliferating and mature chondrocytes); link protein, indian hedgehog (proliferating chondrocytes); PTH-PTHrP receptor (marks chondrocytes at the boundary between the proliferative and hypertrophic zones); and Col10a1 (marks hypertrophic chondrocytes); BMP-6, FGFR3 (marks all chondrocytes). In situ hybridization studies to analyse the biochemical consequences of expressing the transgene at developmental stages 14.5, 16.5 dpc; and postnatal stages: newborn, 5dpp, 5, 10, 15 wks will be performed.

To study possible secondary impact of abnormal bone formation in Col10-13del mutant mice on hematopoiesis we will analyse the long bones in situ for deviations from normal with respect to the morphological appearance of the stroma and haematopoietic cells, incidence of apoptotic and cycling cells and marrow cell number and composition. For this assessment specific molecular markers will be used in situ hybridization studies such as for osteoblasts (Osf2, osteocalcin), osteoclasts (acp5, encoding tartrate resistant acid phosphatase, CSF-1, GM-CSF); chemokines essential for initiation of bone marrow haematopoiesis in ontogeny (e.g. SDF-1); metalloproteinases (BP-1, MMPs) and haematopoietic cytokines (e.g. IL-6, G-CSF) and others.

ii) Further control studies

The following experiments will be performed to determine if the abnormal phenotype is the consequence of:

- a) ectopic expression of the transgene
- b) truncated NC1 domain in collagen X molecules caused by the 13bp del
- c) the novel mutant peptide sequence caused by the 13 bp del which differs from the human
- d) The formation and/or processing and/or matrix deposition of collagen X heterotrimers consisting of mutant and wild-type chains.

To address a) above, transgenic mice will be generated carrying normal Coll10a1 sequences with the same regulatory elements as in the Coll10-13del construct. The phenotype of mice expressing such a transgene will be compared with that of the Coll10-13del mice. The expression pattern of the transgene can be monitored by introducing an internal ribosome entry site sequence followed by an ATG and green fluorescent protein (GFP), at a position downstream of the Coll10a1 translational stop but upstream of the polyA attachment signal.

Alternatively, mice carrying the coll10-13del mutation will be generated using homologous recombination in ES cells to "knock-in" the mutation (Ramires-Solis, R. & Bradley, A Curr. Opin. Biotechnol. 5, 528-533 (1994); Baudoin, C., Goumans, M.-J., Mummery, C. & Sonnenberg. A. Genes & Development 12, 1202-1216).

To address b) a construct containing a SMCD point mutation which results in a premature stop codon in the NC1 domain has been made. This construct will be used to generate transgenic mice and the phenotype of these mice compared with the Coll10-13del mice.

To address c) transgenic mice (Coll10-13delH) will be made carrying the same 13bp deletion mutation except that the

09975607-101101

downstream amino sequence following the 13bp deletion is identical to that in human SMCD mutation. The phenotype of this mouse will be compared with the original Coll10-13del mouse.

5

To address d) transgene mice (Coll10-13del) will be generated on varying genetic backgrounds: heterozygous and homozygous null for Coll10a1. The phenotypes of these mice will be compared with the mice described here, which are Coll10-13del on a wild-type Coll10a1 background. In addition, the biosynthesis and processing of collagen X in chondrocytes isolated from mutant mice will be analyzed.

15

iii) Studying the role of proteoglycans (PGs) and glycosaminoglycans (GAGs) in regulating bone growth.

20

In a recent report, EXT1, a transmembrane glycoprotein present in the ER has been shown to be critical for the expression of cell surface glycosaminoglycans (GAGs). Mutations in EXT1 is the cause of hereditary multiple exostosis (HME), an inherited skeletal disorder characterized by skeletal malformation due to excessive bone growth (32). Type X collagen has been shown to have a strong affinity to bind to proteoglycans (PGs). We propose to test the possibility that the abnormal type X collagen associates differently or fails to associate with proteoglycans (PGs) and/or GAGs in the Coll10-13del mice, thereby causing the bone overgrowth. The

30

biosynthesis of GAGs and PGs will be studied and compared in explant cultures of growth plates from non-transgenic and transgenic mice. In addition the Coll10-13del construct will be transfected into hypertrophic chondrocytes and the effect on PG and GAG biosynthesis studied. These experiments will enable us to determine if PG metabolism has been affected by the abnormal type X collagen.

35

00975607 101101

iv) Cell proliferation and cell cycle regulation in
Coll10-13del hypertrophic chondrocytes

Cell proliferation is controlled by a complicated network
of extracellular and intracellular signalling pathways
5 that process growth regulatory signals and integrate them
into the basic cell-cycle regulatory machinery through
the control of the cyclin-dependant kinases (CDKs). CDKs
are regulated by cyclins (positive regulation) and CDK
inhibitory proteins CKIs (negative regulation). Recently
10 it has been shown that mice lacking the CDK inhibitor
p57kip2 show abnormal proliferation of hypertrophic
chondrocytes and abnormal endochondral ossification
(33,34). The proliferative ability of the hypertrophic
chondrocytes by PCNA (proliferating cell nuclear antigen)
15 immunostaining or BrdU labelling will be studied and
compared. In addition TUNEL (TdT-mediated dUTP-X nick
end labelling assay, Boehringer Mannheim Ltd) assays will
be performed to assess the degree of apoptosis in the
transgenic growth plate compared to non transgenic
20 littermates. Furthermore, PCR differential display will
be used to compare the expression profiles of wild-type
and transgenic growth plates and to clone out genes which
are up-regulated or down-regulated. These studies will
identify key regulators of bone and cartilage growth and
25 the signal transduction pathways which are affected in
the Coll10-13del mice and molecular interactions at the
cartilage-bone junction.

These studies also have implications for understanding
30 the mechanisms underlying the development of bone and
cartilage tumors. Such cartilage and bone tumors include
but are not limited to metastatic bone tumors with
primary tumor arising from the breast and prostate,
osteoid osteoma, osteochondroma, chondroma,
35 osteoblastoma, osteogenic sarcoma, chondrosarcoma,
fibrosarcoma or any other tumors in which increase
cartilage and/or bone formation occurs. In addition,

00975607 101101

these studies will also have implications for understanding the mechanism underlying diseases caused by overgrowth or remodelling problems of bone such as for hyperostosis, exostoses and osteopetrosis, etc.

5

v) Remodelling and turnover of cartilage and bone

The effect of the 13del mutation on the degradation of collagen X will be assessed. To assess degradation of the mutant type X collagen in growth plate, we will isolate the type X collagen substrate from the growth plates of transgenic and wild-type mice (Chan, Azsodi, Fassler and Bateman, unpublished data). Intact and papsinised collagen X will be purified by selective salt fractionation and by affinity chromatography using Sephacryl S-500 resins. This will provide chemical amounts of intact type X collagen (59 kDa) containing both NC1 and NC2 domains as well as pepsined (45 kDa with the NC1 and NC2 domains removed) for metalloproteinase (MMP) cleavage analysis. This will provide data on the susceptibility of the mutant collagen to MMP cleavage. N-terminal amino acid sequencing will be performed on products with major or significant MMP cleavage, MMP cleavage products resolved by SDS-polyacrylamide gel electrophoresis will be sequenced directly following transfer onto PVDF membranes.

10
15
20
25

vi) Bioinformatics and structural analyses

Interspecies comparisons of genes and identification of homologues in vertebrates and invertebrates can reveal insight into gene function and biochemical and genetic pathways. The sequences of genomes of some model organisms have been, or are about to be completely determined. Sequences homologous to the normal and 13-del mutant NC1 domain of collagen X will be screened for using bioinformatics tools in model organisms such as yeast, C. elegans, Drosophila, zebrafish and fugu. Molecular modelling of the normal and mutant NC1 domain

30
35

00075607 101101

may also yield insight into the possible interactions with other molecules.

B. Applied studies for clinical applications

5

There are many different causes of dwarfism. Some are genetic, others environmental or endocrine. Some causes are known, others are as yet undetermined. A number of human conditions can result in a similar phenotype.

10

However, work will need to be undertaken to determine whether they result from the same gene defect.

15

a. The mouse will serve as a useful model for testing the effect of drugs on growth, and the effect of surgical procedures on mice with collagen defects.

20

Drugs like growth hormone has been used to treat achondroplasia (Am J Med Genet 1997 Oct 3;72(1):71-76) with limited effect. The effect of growth hormone in treating other bone dysplasia is not known. A mouse model is an ideal method whereby the effect of growth hormone on skeletal growth can be tested. Additionally, by understanding the basic mechanism by which short stature is produced, new drugs may be devised to stimulate bone growth in these conditions.

25

30

b. Surgical lengthening of bones can increase the height and limb lengths in dwarfs. This has been used to improve on their cosmetic appearance. However, while such bones can lengthen and heal in the lengthened position, what the optimal conditions are to perform the lengthening process is not known. The availability of a mouse model would allow experimentation using an external fixation device to perform lengthening at various rates and frequency. Additionally, by varying the mechanical characteristic of the external fixator the optimal mechanical environment for bone formation can be

35

00075607-101101

ascertained.

5 c. Of particular interest is that the additional bone
formation is manifested by a thickening in the endosteal
and cortical bone. If this process can be made use of in
clinical practice, then it is of advantage over the
currently available bone induction agents (e.g. BMP2 and
BMP7), as these agents result in bone formation in an
uncontrolled manner, resulting in undesirable soft tissue
10 and even muscle calcification.

15 There is the potential to enhance bone formation in
conditions of low bone mass (osteoporosis), and to
improve the quality and speed of bone union after
fracture. This statement is made on the assumption that
the endosteal bone formed is of normal mechanical
characteristic and that the fracture healing occur
uneventfully in these mice. Thus, experiments will be
carried out to determine the mechanical properties of
20 these bones under bending, tension, torsion and
compression. Fracture healing will be assessed in
these mice using an external fixation model, with the
resultant fracture callus assessed mechanically and
histologically. Molecular markers will be used to compare
25 the healing process in Coll10a1-13del mutant and wild-type
mice.

Some possible ways of using this finding to increase bone
mass may be:

- 30 i. Transfect the Coll10a1-13del transgene into
proliferating, hypertrophic chondrocytes and
osteoblasts and assess the effect on in vitro
mineralization using approaches reviewed by Bianco
et al (35).
- 35 ii. To generate the mutant collagen X protein by
recombinant DNA methodology and bacterial/eucaryotic
cell/baculovirus protein expression systems. The

recombinant protein can be implanted in the growth plate or bone and the effect on bone growth and fracture repair assessed. The protein may also be implanted at fracture sites. This may be tested out on normal and osteoporotic bone.

iii. Gene therapy, to deliver the Coll10-13del construct into cells at fracture sites or into osteoblasts in osteoporotic bone.

iv. to complement i)-iii) above, devise and generate vectors for inducible expression of the Coll10-13del gene or the mutant peptide in the NCI domain. These vectors would be derived by modifications of the original Coll10-13del construct by the addition of sequences which may activate expression on addition of an inducer and /or removal of a repressor. Further modifications can also be made to enable switching off of expression of this inducible gene. Such switching off of expression can be achieved by the addition of sequences which can bind repressors of transcription. These vectors may be used to generate transgenic mice in which the timing, site and level of expression of the transgene may be manipulated by induction. Suitable transgenic mice carrying the "inducer" will be generated and crossed to the inducible Coll10-13del mice.

This invention provides different use of the Coll10-13del mice, the Coll10a1 regulatory elements within the vector Coll10-13del, and the novel peptide sequence created by the Coll10a1-13del mutation for basic research into the regulation of bone growth and clinical applications for bone disorders. Specifically, this invention provides the use of the mouse as listed below and the use of the genes which have been affected by expression of the mutant type X collagen. Such genes can be used to develop tools to alter bone formation and growth.

The transgene or key molecules in the pathway revealed as a result of determining the underlying cause of the bone overgrowth can be modified for use in developing a gene or protein based therapy to treat bone and cartilage tumors by reversing or inhibiting chondrocytes proliferation.

The Coll10-13del mouse may be used to:

1. identify key molecules and pathways important for the regulation of bone growth and treatment of dwarfism
- 1.1. To test the effect of drugs or other agents on stimulating bone growth.

Potential drugs for stimulating bone growth in dwarfism can be administered to transgenic and non-transgenic mice which are age and sex matched. The effect of the drug will be assessed by monitoring the animals weight, overall length, and also by measuring the length of the long bones and spine on radiographs taken at regular intervals. In addition, at set time intervals mice will be sacrificed and biochemical and immunohistochemical analyses will be carried out to test for the synthesis of bone-characteristic markers such as type I collagen, osteocalcin, alkaline phosphatase, matrix gla protein, osteonectin, etc.

Groups of control and test animals of different ages and sex will be used to assess the effect of the drug on different age and sex.

Potential drugs (including the use of recombinant human hormones) for increasing bone growth include but is not limited to the natriuretic peptide family of hormones (Yasoda, et al. Natriuretic peptide regulation of endochondral ossification. Evidence for possible roles of C-type natriuretic peptide/guanylyl cyclase-B pathway.

09975607-101101
TOTTOT 20952660

J. Biol. Chem. 1998 273:11695-700), growth hormone (Kidder et al. Effects of growth hormone and low dose estrogen on bone growth and turnover in long bones of hypophysectomized rats. Calcif Tissue Int 1997 61:327-35; Rosen et al., Treatment with growth hormone and IGF-I in growing rats increases bone mineral content but not bone mineral density [published erratum appears in J Bone Miner Res 1995 Nov; 10(11):1836] J Bone Miner Res 1995 10:1352-8.), estrogen (Kidder et al. Calcif Tissue Int 1997 61:327-35), insulin-like growth factor (Kidder et al. Calcif Tissue Int 1997 61:327-35), and parathyroid hormone (Coxam et al. Insulin-like growth factor 1 and parathyroid hormone effects on the growth of fetal rat metatarsal bones cultured in serum-free medium. Biol Neonate 1995 68:368-76) or any combination of the above. Other agents include but are not limited to the expression of a transgene which results in a local delivery of the hormones mentioned above in bone and growth plate (Yasoda, et al. J Biol Chem 1998 273:11695-700), ultrasound, electromagnetic radiation, infrared and laser therapy and acupuncture.

1.2. To test the effect of surgery on increasing bone length.

A limb lengthening device (a modified external fixator that allows for a progressive change in the length of its connecting rods) will be attached to the mouse femur. The femur will then be osteotomised. When early callus formation is observed on radiographs, lengthening of femur will begin (callotaxis). Distraction will be performed under regular radiological monitoring. The frequency and the amount of distraction will be varied to assess the optimal conditions for callotaxis to occur. The animals will be sacrificed at various time points during distraction and at completion of the experiment. The femora will be harvested, and samples sent for mechanical testing and for histological analysis of the

09975607-101101

quality of the new bone formed.

5 A limb lengthening device is a modified external fixator
that allows for a progressive change in the length of its
connecting rods. A number of different systems are in
common use today for increasing bone length in human
subject (Dahl, M.T. and Fischer, D.A. Lower extremity
lengthening by Wagner's method and by callus distraction.
Orthop Clin North Am 1991 22:643-9; Hardy et al. The
10 Sequoia circular fixator for limb lengthening. Orthop
Clin North Am 1991 22:663-75; Price, C.T. and Mann, J.W.
Experience with the Orthofix device for limb lengthening.
Orthop Clin North Am 1991 22:651-61; Saleh, M. and
Burton, M. Leg lengthening: patient selection and
15 management in achondroplasia. Orthop Clin North Am 1991
22:589-99.

2. To test the effect of bone growth stimulating agents
on fracture repair.

20 A fracture repair model in the mouse femur is set up
using an externally fixed method (Andrew, J.G. and
Gregory, J. An externally fixed murine fracture model.
The 25th European symposium on calcified tissues. Bone
25 vol.20, April 1997 p104S; Cheung et al. External fixation
fracture model of the mouse femur. The HKOA annual
congress, 15-16 Nov. 1997).

30 Procedure: The femur will be exposed by an incision on
the lateral aspect of the thigh. The vastus lateralis
muscle is then split to directly expose the mid-shaft of
the femur. The periosteum is not stripped to preserve
the blood supply. The external fxator is applied using
a drill guide to place the four pins. The femur is
35 fractured after drilling holes midway between the pairs
of pins; these holes act as a stress raiser to facilitate
fracture. Skin is closed with an absorbable suture.

000550-10101

The bone growth stimulating agent will be administered to the mice after fracture. A second group of age and sex matched mice will also have the external fixator applied and the bone fractured. No bone growth stimulating agent will be used. This group will act as control.

Outcome will be monitored by regular radiographs, histological and mechanical testing of the harvested one after sacrifice. In addition biochemical, immunohistochemical and gene expression studies will be carried out to assess at the molecular level, whether bone is forming normally.

3. Treatment of human conditions which result in bone over-growth (including osteopetrosis) or mineralization abnormalities by drugs.

Bone overgrowth can occur locally or regionally as a result of certain tumors or in response to certain agents such as physical trauma, ionising radiation, etc. Generalised bone overgrowth can occur in conditions like flurosis (excessive flouride admininstration) and osteopetrosis (cause unknown). In osteopetrosis, it is characterised by a generalized increase in the amount of bone and mineral content of the bone. Both the cortical and trabecular bone are affected with a resultant reduction in the marrow spaces within the medullary cavity. The cause of the condition is not known. The mouse may be used to:

- i. Identify key molecules or pathways important for the regulation of bone growth and bone overgrowth.
- ii. To make drugs which can be used to reverse or inhibit this process of bone overgrowth.
- iii. Act as a model to test the effect of other drugs on reversing or inhibiting bone overgrowth.

Use of the model to identify drugs for treatment of bone overgrowth

5 By understanding the mechanism by which the Col10-13del transgene can cause bone overgrowth, and the regulatory pathways involved, a drug may be devised which can act as an inhibitor of bone growth for example one which inhibits osteoblast proliferation and/or differentiation. The drug may also be an antagonist to some key step in
10 the regulatory pathway.

Use of the model to test the effect of drugs on bone overgrowth

15 The method involves administration of the drug to a group of transgenic animals, while another group of age and sex matched transgenic will be given a placebo to act as control. The effect of the drug on bone density will be assessed by regular radiological monitoring.
20 Additionally, bone densitometry, micro-CT analysis or other more sensitive methods may be used in future to assess the effect of the drug on bone density. At regular time intervals the mice will be sacrificed and the bones examined histologically. Bone density may also
25 be estimated using electron back-scattering analyses combined with electron microscopy. Histomorphometric analysis will be carried out to quantify the effect on the bone content. Biochemical and immunohistochemical analyses will also be carried out to study the level of
30 synthesis of bone-characteristic markers such as type I collagen, osteocalcin, alkaline phosphatase, matrix gla protein, osteonectin, etc.

4. The transgene can be used or modified to be used as a method of regulating bone growth and switched on
35 in conditions where more bone is required. For example - improving the amount of bone in patients with osteoporosis and increasing the rate of fracture healing using a gene therapy approach which may include an inducible/switchable approach as

09975607 101101
TOTTOT 20952660

described above.

5. The mouse can be used to find potentially new genes and pathways responsible for regulating bone growth with implications for human bone disorders
6. If a human inherited condition caused by a regulatory defect in COL10A1 is found, the use of the mutation for diagnosis.
7. The use of the regulatory sequences in the transgene Col10-13del to target expression of therapeutic compounds to the growth plate by transgenesis.
8. The use of the novel peptide sequence in the NC1 domain created by the 13bp deletion and antibodies raised against this sequence, if it should turn out to play a role in causing the abnormal bone growth.

Identification of genetic modifiers of bone growth

Since the severity of phenotype in the Col10-13del mice varied within a line, and these mice are essentially a combination of 129sv/J, CBA and C57BL6 backgrounds, these mice will also serve as models to study the influence of genetic background on bone growth. Mice expressing the mutation on different bred background may be generated by breeding these mice to generate congenic mutant mice with different inbred background. Comparison of the phenotypic severity and extent of cartilage and bone growth will shed insight into the degree to which the bone overgrowth is affected by genetic background. Should differences in the bone overgrowth found in different inbred backgrounds, the mice could be used to identify genetic loci which account for genetic variation in bone growth. These loci may be detected using genetics, linkage analyses and highly polymorphic markers such as single nucleotide polymorphism (SNPs).

Molecular consequences of expressing the mutant (Col10-12del)protein in hypertrophic chondrocytes

As discussed in section number iv above, the expression of the cell cycle regulator p57kip2 in the growth plates of non-transgenic and Coll10-12del mice has been analyzed using immunohistochemistry (antibodies, gift of Dr. Anne Fergusson-smith, Dept of Anatomy, Cambridge University, UK). In non-transgenic controls, strong staining for p57kip2 was seen in the hypertrophic chondrocytes throughout the hypertrophic zone. However, the immunostaining showed areas of reduced staining within the hypertrophic zone of Coll10-13del mice. Using antibodies to POCNA (Boehringer Mannheim), strong immunostaining was seen in hypertrophic chondrocytes of the Coll10-13del mice which was not seen in the non-transgenic controls. These data are consistent with abnormal proliferation of the hypertrophic chondrocytes in the Coll10-13del mice which may in part account for the expansion of the hypertrophic zone in these mice.

It has been suggested, as discussed in section number iv above, that there may be a problem with secreting the mutant collagen X protein in the transgenic mice (see experimental details above). Antibodies to the altered peptide sequence arising from the 13 del mutations have been generated (see experimental details in section number i above). This antibody has been used to stain growth plates sections from the Coll10-13del and control mice. Immunostaining using antibodies specific to the mutant sequence in the Coll10-13del protein shows strong intracellularly and almost undetectable signal extracellularly. No staining was seen in non-transgenic samples. Confirming the transgene specificity of the antibody. This data is consistent with an impaired ability to secrete the mutant protein. By contrast, immunostaining with antibodies to collagen X (gift of Bjorn Olsen, Harvard University) which would stain both the endogenous normal and mutant protein, showed an extracellular pattern of expression. The cells in the

00075607-101101

trabeculae may be hypertrophic chondrocytes in the cartilaginous remnants within trabeculae, or osteoblasts.

5 There was also increased staining of hypertrophic
chondrocytes in the growth plate and cells in the
trabeculae of Col10-13del mice using antibodies to heat
shock protein 47 (hsp47), a chaperone of collagens (Ref:
Expression K Matrix Biol 1998: 7 379-386). There was
10 little staining with hsp47 in the hypertrophic
chondrocytes of nontransgenic mice. Taken with the
electron microscopy data showing the presence of engorged
endoplasmic reticulum, this observation would be
consistent with intracellular accumulation of protein due
to difficulty with secretion and support hypothesis (see
15 section f above).

Active degradation of intraceullular proteins can have
significant implication on cellular behavior. Interaction
of unfolded X chains with chaperones such as protein
20 disulfide isomerase and Hsp47, glcoprotein-specific
chaperones such as calnexin and calreticulin, BiP and
Ubiquitin (see Ellis RJ Molecular chaperones: pathways
and networks, Curr Biol 1999 9:4 R137-139) will also be
assessed in further studies by immuno-coprecipitation
25 methods. This will give insights into the intracellular
handling of unassembled collagen X chains and preferred
pathway of clearance.

Expression has been seen (by in situ hybridization and by
30 immunostaining) of the Col10-13del transgene in cells
which are likely to be osteoblasts in the cortical bone.
This finding indicates that expression of the mutant
protein in both bone and hypertrophic cartilage may be
the cause of the altered phenotype in Col10-13del mice.

35

In collaboration with Dr. Davine Opelstein (Biochemistry
Dept, University of Hong Kong) the composition of the

09975607 101101

marrow cells in wild-type and Col10-13del transgenic mice has been analyzed. It was found that at 26 weeks of age, a marked increase in the numbers of adipocytes. There also appeared to be an increased number of blood vessels in the bone. There appeared to be an increase in the volume of the marrow cavity, and an abundance of total haematopoietic cells in wildtype compared to transgenic mice.

Consistent with an effect on haemetopoiesis was observation of abnormal morphology of the thymus in one of the transgenic mice. Further mice will need to be analyzed to determine if this is a consistent finding. These changes indicate an effect of the altered microenvironment of the transgenic bone on the development of cells within the marrow cavity. It is unknown if this is a direct effect or an indirect consequence whereby haematopoeitic cells remain constant by adipocytes are increased to compensate for the increased size of the marrow cavity in the transgenic mice. The mice are also closely monitored for signs of srthritis, such as rheumatoid arthritis or osteoarthritis).

25

0995607-101101

References

1. Mayne, R., and Irwin., M.H. (1986) in Articular cartilage biochemistry (Kuettner, K.E., Schleyerbach, R., and Hascall, V.C., eds) pp. 23-35, Raven Press, New York.
5
2. Schmid, T.M., and Linsenmayer, T.F. (1987) in Structure and Function of Collagen Types (Mayne, R., and Burgeson, R.E., eds) pp. 223-259, Academic Press, Orlando, Florida 20.
10
3. Kong, R.Y.C., Kwan, K.M., Lau, E.T., Thomas, J.T., Boot-Handford, R.P., Grant, M.E., and Cheah, K.S.E. (1993) Eur. J. Biochem. 213, 99-111.
15
4. Schmid, T.M., and Linsenmayer, T.F. (1990) Dev. Biol. 138, 53-62.
5. Kwan, A.P.L., Cummings, C.E., Chapman, J.A., and Grant, M.E. (1991) J. Cell Biol. 14, 597-604.
20
6. Kirsch, T., and Pfaffle, M. (1992) FEBS Lett. 310, 143-147.
7. Anderson, H.C. (1989) Lab. Invest. 60, 320-330.
25
8. Kirsch, T., Swoboda, B., and Von der Mark, K. (1992) Differentiation 52, 89-100.
9. Gibson, G.J., Bearman, C.H., and Flint, M.H. (1986) Coll. Rel. Res. 6, 163-184.
30
10. Schmid, T.M., and Linsenmayer, T.F. (1985) J. Cell Biol. 100, 598-605.
11. Bonen, D.K., and Schmid, T.M. (1991) J. Cell Biol. 115, 1171-1178.
35

09975507-104101
FOTOT 20952660

12. Schmid, T.M., Bonen, D.K., Luchene, L., and
Linsenmayer, T.F. (1991) *In Vivo* 5, 533-540.
13. Poole, A.R., and Pidoux, I. (1989) *J. Cell Biol.*
109, 2547-2554.
14. Schmid, T.M., Popp, R.G., and Linsenmayer, T.F.
(1990) *Ann. NY Acad. Sci.* 580, 64-73.
15. Warman, M.L., Abbott, M., Apte, S.S., Hefferon, T.,
McIntosh, I., Cohn, D.H., Hecht, J.T., Olsen, B.R.,
and Francomano, C.A. (1993) *Nature Genet.* 5, 79-82.
16. Wallis, G.A., Rash, B., Sweetman, W.A., Thomas,
J.T., Super, M., Evans, G., Grant, M.E., and
Boot-Handford, R.P. (1994) *Am. J. Hum. Genet.*
54, 169-178.
17. Wallis, G.A. (1993) *Curr. Biol.* 3, 687-689.
18. McIntosh, I., Abbott, M.H., and Francomano, C.A.
(1995) *Hum. Mutat.* 5, 121-125.
19. Lachman, R.S., Rimoin, D.L., and Spranger, J. (1988)
Pediatr. Radiol. 18, 93-102.
20. Horton, W.A., and Hecht, J.T. (1993) in *Connective
tissue and its heritable disorders, molecular
genetic and medical aspects* (Royce, P.M., and
Steinmann, B., eds) pp. 641-675, Wiley-Liss, New
York.
21. Jacenko, O., LuValle, P.A., and Olsen, B.R. (1993)
Nature 365, 56-61.
22. Jacenko, O., LuValle, P., Solum, K., and Olsen, B.R.
(1993) *Prog. Clin. Biol. Res.* 383B, 427-436.

00952660 10101

23. Rosati, R., Horan, G.S.B., Pinero, G.J., Garofalo, S., Keene, D.R., Horton, W.A., De Crombrughe, B., and Behringer, R.R. (1994) Nature Genet. 8, 129-135.
- 5 24. K. M. Kwan, M.K.M. Pang, S. Zhou, S.K. Cowan, R.Y.C. Kong, T. Pfordte, B.R. Olsen, D. Sillence, P.P.L. Tam, & K.S.E. Cheah (1997) J. Cell Biol. 136 459-471.
- 10 25. Chan, D., Jacenko, O. (1998) Matrix Biol. 17:169-184.
26. Chan, D., Cole, W.G., Rogers, J.G., and Bateman, J.F. (1995) J. Biol. Chem. 270, 4558-4562.
- 15 27. Chan, D., Weng, Y.M., Hocking, A.M., Golub, S., McQuillan, D.J., and Bateman, J.F. (1996) J. Biol. Chem. 271, 13566-13572.
- 20 28. Horton, R.M., Hunt, H.D., Ho, S.N., Pullen, J.K., and Pease, L.R. (1989). Gene, 77:61-68.
29. Elima, K., Eerola, I., Rosati, R., Mets=E4ranta, M., Garofalo, S., Perala, M., de Crombrughe, B., and Vuorio, E. (1993) Biochem. J. 289:247-253.
- 25 30. Cheah, K.S.E., Levy, A., Trainor, P.A., Wai, A.W.K., Kuffner, T., So, C.L., Leung, K.K.H., Lovell-Badge, R.H., and Tam, P.P.L. (1995) J. Cell Biol. 128, 223-237.
- 30 31. Vortkamp, A., Lee, K., Lanske, B., Segre, G.V., Kronenberg, H.M., and Tabin, C.J. (1996) Science 273, 613-622.
- 35 32. McCormick, C., Leduc, Y., Martindale, D., Mattison, K., esford, L.E., Dyer, A.P., and Tufaro, F. (1998)

TDIOT 20952660

Nature Genetics 19:158-161.

33. Zhang, P., Liegeois, N.J., Wong, C., Finegold, M., Hou, H., Thompson, J.C., Silverman, A., Harper, J. W., DePinho, R.A., Elledge, S.J. (1997) Nature 387:151-158.
34. Yan, Y. Frisen, J., Lee, M-H., Massague, J., Barbacid, M. (1997) Genes & Dev. 11: 973-983.
35. Bianco, P., Cancedda, F.D., Riminucci, M., Cancedda, R. (1998) Matrix Biol. 17:185-192.
36. Nagata, K., (1998) Matrix Biol. 7:379-386.
37. Ellis, R.J., (1999) Curr. Biol. 9:4 R137-139.